



## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)


(51) International Patent Classification <sup>6</sup> : <b>G01N 33/50, 33/574</b>		<b>A1</b>	(11) International Publication Number: <b>WO 99/08108</b>
			(43) International Publication Date: 18 February 1999 (18.02.99)
(21) International Application Number: PCT/NL98/00457		(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, HR, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).	
(22) International Filing Date: 11 August 1998 (11.08.98)			
(30) Priority Data: 97202501.9 12 August 1997 (12.08.97) EP			
(71) Applicant (for all designated States except US): LEADD B.V. [NL/NL]; Wassenaarseweg 72, NL-2333 AL Leiden (NL).			
(72) Inventor; and			
(75) Inventor/Applicant (for US only): NOTEBORN, Mathieu, Hubertus, Maria [NL/NL]; Sternstraat 15, NL-2352 EH Leiderdorp (NL).			
(74) Agent: SMULDERS, Th., A., H., J.; Vereenigde Octrooibureaux, Nieuwe Parklaan 97, NL-2587 BN The Hague (NL).			
		Published With international search report.	

(54) Title: DETERMINING THE TRANSFORMING CAPABILITY OF AGENTS

## (57) Abstract

The invention relates to activation of apoptin-induced apoptosis by different cell-transforming agents in normal and/or cancer-prone cells. Apoptin or also called VP3 is a viral protein derived from the Chicken anemia virus. Also the invention relates to preventive anti-tumor therapies of normal and/or cancer-prone cells. Treatment of normal and/or cancer-prone cells with tumor-inducing agents will activate apoptin-induced apoptosis, resulting in the elimination of potential tumor cells. Also the invention relates to diagnosis of cancer agents. Agents with tumor activity can be examined by expressing them in normal cells and analyzing their capability of enabling apoptin-induced apoptosis.

## Overview VP3 transgenic mice



**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon	KR	Republic of Korea	PL	Poland		
CN	China	KZ	Kazakstan	PT	Portugal		
CU	Cuba	LC	Saint Lucia	RO	Romania		
CZ	Czech Republic	LI	Liechtenstein	RU	Russian Federation		
DE	Germany	LK	Sri Lanka	SD	Sudan		
DK	Denmark	LR	Liberia	SE	Sweden		
EE	Estonia			SG	Singapore		

Title: DETERMINING THE TRANSFORMING CAPABILITY OF AGENTS

The present invention relates to the field of cancer diagnosis and treatment, as well as to the field of determining the transforming capability of suspected tumorigenic or tumor promoting agents (the two terms will be  
5 used interchangeably herein).

The common denominator of the present invention is that all of the above fields are fields in which apoptin or derivatives and/or fragments thereof (hereinafter all referred to as apoptin or apoptin-like activity) can be  
10 applied according to the invention. Apoptin is a protein originally found in Chicken Anemia Virus (Noteborn et al., 1991) and was originally called VP3. The apoptotic activity of this protein was discovered by the group of the present inventors (Noteborn et al., 1994).

15 As stated above the present invention makes use of the apoptosis inducing effect of apoptin.

Apoptosis is an active and programmed physiological process for eliminating superfluous, excessively damaged or malignant cells (Earnshaw, 1995, Duke et al., 1996).

20 Apoptosis is characterized by shrinkage of cells, segmentation of the nucleus and fragmentation of the cytoplasm, condensation and cleavage of DNA into domain-sized fragments, in most cases followed by internucleosomal degradation. The apoptotic cells become fragmented into  
25 membrane-enclosed apoptotic bodies. Finally, neighbouring cells and/or macrophages will rapidly phagocytose these dying cells (Wyllie et al., 1980, White, 1996). Cells grown under tissue-culture conditions and cells from tissues can be analysed for signs of apoptosis with agents staining  
30 chromosomal DNA, as e.g. DAPI or propidium iodide, which stains normal DNA (chromatic) strongly and regularly, but

apoptotic chromatin weakly and/or irregularly (Noteborn et al., 1994, Telford et al., 1992).

The apoptotic process can be initiated by a variety of regulatory stimuli (Wyllie, 1995, White 1996, Levine, 1997).

5 Changes in cell survival rate play an important role in human pathogenesis, e.g. in cancer development, which is caused by enhanced proliferation and/or by decreased cell death (Kerr et al., 1994, Paulovich, 1997). A variety of chemotherapeutic agents and radiation have been demonstrated to induce  
10 apoptosis in tumor cells which, in many instances is mediated via the tumor suppressor protein p53 (Thompson, 1995, Bellamy et al., 1995, Steller, 1995, McDonnell et al., 1995).

Many tumors, however, acquire a mutation in p53 during their development, often correlating with poor response to cancer  
15 therapy. Transforming proteins of DNA tumor viruses inactivate p53 indirectly or by directly binding to it (Teodoro, 1997). An example of such an agent is the large T-antigen of the DNA tumor virus SV40. In certain hemopoietic tumors, a high expression level of the Bcl-2-oncogene is  
20 associated with a strong resistance to various apoptosis-inducing chemotherapeutic agents (Hockenberry 1994, Sachs and Lotem, 1997). For such cancers, that are resistant to many cytotoxic agents, alternative anti-tumor therapies are under development based on induction of apoptosis (Thompson, 1995  
25 and Paulovich et al., 1997).

Apoptin is a small protein derived from the chicken anemia virus (CAV; Noteborn and De Boer, 1995, Noteborn et al., 1991, Noteborn et al., 1994), which can induce apoptosis in human malignant and transformed cell lines, but not in  
30 untransformed diploid human cells. In vitro, apoptin fails to induce programmed cell death in normal lymphoid, dermal fibroblastic, epidermal, endothelial and smooth-muscle cells. However, when normal cells are transformed e.g. by the transforming genes of SV40, they become susceptible to  
35 apoptosis by apoptin. (Danen-van Ooschot, 1997 and Noteborn, 1996). Long-term expression of apoptin in normal human

fibroblasts revealed that apoptin has no toxic or transforming activity in these cells. In normal cells, apoptin was found to be localized predominantly in the cytoplasm, whereas in transformed or malignant cells, it was  
5 located in the nucleus, suggesting that the localization of apoptin is related to its death-inducing activity (Danen-van Oorschot et al. 1997).

Furthermore, we have established that apoptin can induce apoptosis in the absence of functional p53 (Zhuang et al.,  
10 1995a), and cannot be inhibited by Bcl-2, Bcr-abl (Zhuang et al., 1995), the Bcl-2-associating protein BAG-1 and the caspase-inhibitor cow-pox protein CrmA (Danen-Van Oorschot, 1997, Noteborn, 1996). Finally, it appears that cells that are only immortalized and thus minimally transformed, can  
15 also be killed by apoptin.

Therefore, apoptin is an extremely potent anti-tumor agent, also for tumors that are not or less susceptible to (chemo)therapeutic agents due to the lack of functional p53, (over)-expression of Bcl-2 or other apoptosis-inhibiting  
20 genes. The fact that apoptin does not induce apoptosis in normal human cells, suggests that a toxic effect of apoptin treatment in vivo will be very low. In addition, it appears, that even premalignant, minimally transformed cells, may be sensitive to the death-inducing effect of apoptin.

25 Knowing that apoptin is quite safe in normal cells, but that as soon as a cell becomes transformed and/or immortalized (the terms may be used interchangeably herein) the present inventors designed some uses based on this finding.

Thus the invention provides a method for determining the  
30 transforming capability of a possible transforming agent, comprising providing a non-transformed cell with inducible apoptin-like apoptotic activity, exposing said cell to said transforming agent and determining the localization of said apoptotic activity within said cell or determining the  
35 induction of apoptosis in said cell.

It is to be understood that in the above apoptotic activity also refers to the entity having said activity.

It is preferred to provide said cell with said apoptotic activity by transducing said cell with a recombinant nucleic acid molecule encoding said activity. Apoptin-like activity is herein defined as any (preferably proteinaceous) substance having similar activity as VP3 or apoptin of chicken anemia virus. Specifically included in said definition are allelic variants, derivatives and/or fragments of apoptin, wherein derivatives are defined as having amino acid replacements which do not result in the loss of all apoptotic activity. It is to be understood that similar activity means that the kind of activity is the same although the amount may differ. The methods according to the invention are especially suitable in applications whereby said possible transforming agent is a proteinaceous substance. This allows for said proteinaceous substance to be co-expressed in said non-transformed cell with said apoptotic activity. The exemplified proteinaceous substance is the large T-antigen of SV40 or a functional equivalent thereof.

The invention also provides modifications on the Apoptin gene resulting in changes on the apoptin protein enabling apoptin to enter the nucleus in non-transformed and transformed/tumorigenic cells, resulting in the induction of apoptosis. The apoptin protein is enlarged with a nuclear localization signal of SV40. Specifically included in said definition of apoptin are allelic variants, derivatives and/or fragments of apoptin, wherein derivatives are defined as having amino acid replacements which do not result in the loss of all apoptotic activity. This allows for the apoptin protein to be expressed in non-transformed cells with said apoptotic activity. Apoptin fragments with said apoptotic activity but not able to enter the nucleus of non-transformed or transformed cells by its own sequences, are able to enter the nucleus by means of the modifications and induce apoptosis.

The invention further provides a method for determining the predisposition of a cell to become a tumor cell, by providing said cell with inducible apoptin-like apoptotic activity and subjecting said cell to relatively mild tumorigenic activity and determining apoptosis in said cell and/or determining the localization of said apoptotic activity in said cell. In this case the suspected transforming agent as discussed hereinbefore is already present in said cell as a mutation leading to oncogenic or tumorigenic activity. In that case the fact that apoptin only induces apoptosis in cells which have already been transformed leads to the possibility to check whether cells have a mutation which will lead to immortalization or transformation upon mild exposure to transforming activity, such as UV-irradiation and X-ray treatment.

In this manner the likelihood of a set of cells to lead to cancer can be determined. This of course leads to applications in the field of diagnosis of the chances of people having a hereditary risk of getting cancer and giving preventive treatment, which is another object of the present invention. This kind of diagnosis can also be applied in advising people of the likelihood of their children to be predisposed for cancer.

Thus the invention also provides a method for determining the predisposition of a subject for hereditary types of cancer, comprising subjecting a sample of a relevant subset of cells of such a subject to a method as disclosed herein. And the invention further provides a method for determining a gene mutation having oncogenic and/or transforming activity in a cell comprising subjecting said cell to a method according to the invention.

As stated hereinbefore it is another object of the present invention to provide means for the prophylactic treatment of subsets of cells in a person, which subset of cells is cancer-prone. These means include a nucleic acid encoding

apoptin-like activity, preferably provided in the form of a gene delivery vehicle. Gene delivery vehicles can be of viral origin or other. Many vehicles have been disclosed in the art and are known to the skilled person. They include but are not limited to adenoviral vectors, preferably in the form of adenoviral particles; retroviral vectors, preferably as recombinant retroviruses; the same kind of vectors but derived from other viruses; liposomes or other carrier molecules, etc.

10 The invention also provides a diagnostic test kit for carrying out a method according to the invention in determining the tumorigenic capability of an agent, comprising a non-transformed cell transduced with a nucleic acid encoding apoptin or a functional derivative or fragment thereof and optionally all other material necessary to carry out the test and detecting the result.

The invention further provides a diagnostic test kit for carrying out a method according to the invention for determining the cancer-proneness of cells, comprising a nucleic acid encoding apoptin or a functional derivative or fragment thereof capable of transducing a eukaryotic cell and capable of being expressed in such a cell and optionally all other material necessary to carry out the test and detecting the result. Preferably a means for subjecting a cell to a mild tumorigenic activity, such as UV-irradiation and X-ray treatment is also provided.

The invention also provides a method for studying the induction of Apoptin-induced apoptosis by transforming agents such as chemicals, viruses, UV- and X-irradiation in a transgenic mouse model, which results in inhibition of tumor formation. The transgenic-apoptin mice can be used for analysing the anti-tumor effect of Apoptin in transgenic chimeras carrying hereditary types of cancer and able to express Apoptin. Furthermore, the effect of the expression of apoptin in an in-vivo model can be studied by means of the described transgenic-apoptin mice.



**Detailed description of the invention.**

We have previously shown that the viral protein apoptin induces apoptosis in cultured transformed cells of both human  
5 and rodent origin, but not in normal human cells. We have now observed that apoptin fails to induce apoptosis in cultured murine (or rat) embryonic fibroblasts. (the cultures were derived from 16-18 days old mouse (rat) embryos.) This shows that apoptin may also be expressed in the intact embryo  
10 without causing toxicity, at least embryos of a not too early stage of embryonic development.

We have now been able to produce transgenic-apoptin mice, which are viable. We provide evidence that constitutive expression of apoptin in a transgenic mouse does not result  
15 in lethal or other life-threatening/life-limiting effects. We have also observed that co-transfection of cultured normal human fibroblasts with the apoptin gene and the SV40 transforming genes will activate the apoptotic process, which is accompanied by translocation of the apoptin protein from  
20 the cytoplasm, where it accumulates initially, to the nucleus.

The described invention provides the basis for amino-acid additions to the apoptin protein or its fragments enabling those to enter and/or accumulate in the cellular  
25 nucleus, resulting in the induction of apoptosis.

The invention provides the basis for a diagnostic test for detection of potentially transforming genes. Normal diploid mammalian cells, such as human and/or rodent cells are used for such a test. To that end, the normal diploid  
30 cells are co-transfected with a plasmid containing the gene(s) to be studied and the plasmid encoding apoptin, or transfected with the gene(s) to be studied and infected with a viral vector expressing apoptin. Induction of apoptin-induced apoptosis and/or presence of apoptin in the nucleus  
35 show that the examined gene harbors transforming/tumorigenic potential.

Furthermore, we have discovered that diploid human cells isolated from individuals who carry a germ line mutation in a tumor-suppressor gene, and as a result are predisposed to develop a certain spectrum of tumors (herein also referred to as cancer-prone), are resistant to the apoptosis-inducing effect of apoptin, just as diploid cells from healthy individuals, however, become sensitive if the cell cultures are irradiated with ultraviolet light.

This allowed us to develop a diagnostic assay for predicting cancer-proneness. In families with a hereditary predisposition to cancer due to a germ-line mutation in a tumor suppressor gene, it is often not possible, without an extensive analysis of chromosomal DNA, to predict whether a family member is afflicted and carries the disease gene. Our results show that this is in a simple way, by making use of the apoptin gene. To that end, diploid skin fibroblasts or lymphocytes are isolated from the individual to be tested, and the cultured cells are transfected with the apoptin gene, followed by irradiation with UV-light (266nm). If the transfected cells become apoptotic after UV irradiation but fail to enter apoptosis without UV exposure, then this is a (strong) indication that the individual is cancer-prone. Not all types of cancer predisposition, that are due to a mutation in a tumor suppressor gene, have been tested with the Apoptin/ UV assay as yet. There is however no reason to assume that the same phenomenon will not be observed in other cancer-prone cells. A similar diagnostic test for predicting cancer-proneness can also be carried out by using X-ray treatment instead of UV irradiation.

The invention also allows us to obtain more information on the molecular basis of cancer-proneness and its relationship with certain stress-responses, such as Enhanced Reactivation (ER) (Abrahams et al., 1996). Enhanced Reactivation is one of the reactions of normal (human) cells to certain DNA-damaging agents, and appears to reflect the cells' susceptibility to oncogenic transformation.

The invention will be explained in more detail in the following experimental part. This only serves for the purpose of illustration and should not be interpreted as a limitation of the scope of protection.

5

### Experimental part

#### **Cells and cell culture conditions**

Rat embryo fibroblasts (REF) were prepared from 14-day-old rat embryos. The cells were thawed from liquid nitrogen, cultured in DMEM supplemented with 10% fetal calf serum, and transfected with plasmid DNA at cell passage 2.

Mouse embryo fibroblasts (MEF) were prepared from p53+/+ mice or from p53 -/- knock-out mice (Tyler Jacks, 1994 and 1996; Tyler Jacks et al., 1994). The cells were grown on Corning dishes in F15 medium supplemented with 10% fetal calf serum. P19 cells are derived from a mouse embryonal carcinoma/teratocarcinoma (Burney et al., 1982). The cells were grown on gelatinized Petri dishes in DMEM supplemented with 8% fetal calf serum.

BRK/xho cells are prepared from baby rat kidney cells, by transforming with the adenovirus type 5 E1 region (Schrier et al., 1983). The cells were cultured in DMEM supplemented with 10% fetal calf serum.

Human diploid foreskin fibroblasts VH10 and VH25 (Klein et al., 1990) were grown in Dulbecco's modified Eagle medium (DMEM) supplemented with 10% fetal-calf serum.

Primary cultures of human epidermal keratinocytes (FSK-1) were initiated in complete medium as described (Rheinwald and Green, 1975), with minor modifications according to M. Poncet et al., 1981, and cultured in keratinocyte serum-free medium (KSFM) afterwards. For the experiments described here, passage number 3 was used.

F9605 cells are diploid fibroblasts which are p16-/-, derived from patients with dysplastic nevus syndrome (DNS), which is postulated to be a precursor of familial atypical multiple

mole-melanoma (FAMMM) syndrome (Gruis et al., 1995). The cells were grown in DMEM with 10% fetal calf serum.

GM1492 cells are human diploid fibroblasts, which do not express p53 and are derived from patients with Bloom's

5 Syndrome, an autosomal recessive disorder with a high cancer incidence (Van Laar et al., 1994) The cells were grown in DMEM containing 10% fetal calf serum.

LF2675T are diploid skin fibroblasts from patients with Li-Fraumeni syndrome, (LFS). This disease is characterized by a

10 germline mutation in one allele of the p53 gene and an early onset of various types of cancer (Srivastava et al., 1990; Abrahams et al., 1996). The cells were grown in DMEM supplemented with 10% fetal calf serum.

401 cells are diploid skin fibroblasts from individual of

15 Lynch type 2 syndrome family with a high incidence of various types of cancer. The cells were derived from an individual who died of breast cancer (Abrahams et al., 1996). The cells were grown in DMEM supplemented with 10% fetal calf serum.

All culture media were obtained from GIBCO/BRL and contained

20 the antibiotics penicillin and streptomycin.

#### **Irradiation of the cell cultures**

The conditioned medium was removed from the cultures and the cells were rinsed twice with PBS. After removal of the

25 PBS, the cultures were UV-irradiated as described previously (Abrahams et al., 1984), or treated with X-rays (5 gray) by

using an Andrex 225 SMART (Andrex St, Copenhagen) at 200 KV, 4 mA with a 1-mm Al filter. Dose and dose rate were monitored

with a PTW dosimeter. After UV treatment, conditioned medium

30 was returned, and the cultures were incubated at 37°C.

### Plasmids

The expression plasmid pCMV-VP3 contains the CAV DNA sequence encoding the apoptin protein exclusively (nt 427-868; Noteborn et al., 1991, Noteborn and De Boer, 1996), and  
5 plasmid pCMV-des encodes desmin, a structural protein of muscle cells (Menke et al. 1997). Plasmid pCMV-neo is used as the "empty" negative control for plasmids encoding gene products with a potential effect on the apoptin-induced apoptosis. All expressed genes are under the regulation of  
10 the cytomegalovirus (early) enhanced promoter. Plasmid SV40 contains the SV40 origin-defective (ori-) early-region clone including both SV40 large T antigen- and small T antigen-coding regions regulated by their own promoter (Dinsart et al., 1984). The plasmid pR-s884 expresses a  
15 complete SV40 large T antigen and a truncated small T antigen under the transcriptional control of the long terminal repeat (LTR) of Rous sarcoma virus (RSV; De Ronde et al., 1989; Smits et al., 1992). The plasmid PR-SVt contains cDNA sequences coding for the SV40 small T gene that was fused to  
20 the RSV LTR (Philips and Bundell, 1988).

The expression plasmid 21EcoA, which consists of the murine H-2Kb histocompatibility antigen gene/promoter regions (Mellor et al., 1982) and pBr327 sequences is a gift from  
25 Prof Dr Frank Grosveld, Erasmus University, Rotterdam, The Netherlands. The plasmid 21EcoA contains a NotI within the first exon of the H-2Kb gene, which enables the integration of a foreign gene becoming regulated by the H-2Kb promoter. The BamHI fragment containing the sequences encoding apoptin was isolated from pCMV-VP3 and cloned in the NotI site of  
30 21EcoA plasmid by using NotI-BamHI linkers. The final plasmid containing the apoptin gene under the regulation of the H-2Kb promoter was called p21EcoA-VP3. Subsequently, the EcoRI-site next to the apoptin gene was deleted by linearization of plasmid p21EcoA-VP3 at this specific EcoRI site, and  
35 treatment with Klenow polymerase treatment. The plasmid is called p21EcoA-Vp3-Eco.

The DNA fragment containing the H-2kB expression unit with the apoptin gene was separated from the procaryotic DNA sequences by means of EcoRI digestion and agarose-gel electrophoresis.

- 5 Plasmid DNA was purified by centrifugation in a CsCl gradient and column chromatography in Sephacryl S500 (Pharmacia, Sweden).

#### **Transient Transfection**

- 10 The cells were transfected in monolayed cultures with transfection agent DOTAP (Boehringer, Mannheim, FRG) essentially as described by Fischer et al., 1996, or transfected with plasmid DNA by calcium-phosphate precipitation as described by Graham and Van der Eb (1973).

15

#### **Indirect immunofluorescence.**

- All cells were grown on glass microscope slides. The slides were either uncoated (VH10, VH25), or coated with 3-amino-propyltriethoxysilane (TESPA; FSK-1). The cells were  
20 fixed with 80% acetone for 10 min at room temperature, and used for indirect immunofluorescence as described (Van den Heuvel, 1990). To demonstrate the presence and/or cellular localization of apoptin in transfected cells, mouse monoclonal antibody (Mab) CVI-CAV-85.1 (85.1; Noteborn et al., 1991); for human desmin the mouse Mab 33 (Monosan, Uden, The Netherlands); for SV40 T antigens Pab 419, kindly provided by Dr A.-G Jochemsen, University of Leiden, The Netherlands, was used. Fluorescein isothiocyanate-labeled goat anti-mouse antibody (Jackson Immunoresearch Laboratories  
25 Inc., West grove PA, USA) was used as second antibody. Nuclear DNA was stained with 2,4-diamino-2-phenylindole (DAPI).

#### **The generation of transgenic-apoptin mice.**

- 35 For the generation of transgenic-apoptin mice, fertilized oocytes from the FVB mouse strain and mice from

the murine strain were used as fosters. The micro-injections were carried out in the male pronuclei according to Brinster et al. (1981). Per micro-injection 500 copies of the EcoR1 DNA fragment, derived from plasmid p21EcoA-Vp3-Eco, containing the required H-2Kb transcription unit and the complete apoptin gene was injected.

## Results and Discussion

### Apoptin induces apoptosis in rodent transformed cells but not in normal embryonic cells.

To examine whether apoptin fails to induce apoptosis in normal embryonic rodent cells, cultures of mouse embryo cells and rat embryo cells were transiently transfected with a plasmid encoding apoptin. As a negative control, cells were transfected with a plasmid encoding desmin, which does not have apoptotic activity. Cells expressing apoptin were screened via indirect immunofluorescence with Mab 85.1, and cells expressing desmin with mouse Mab 33. Induction of apoptosis in apoptin- or desmin-positive cells was analyzed with the help of DAPI, which causes a regular staining in intact nuclei, but an irregular and/or weak staining in apoptotic nuclei.

Five days after transfection around 10-20% of the desmin-positive cells were apoptotic, which is the basal level most likely due to the transfection event (data not shown, Menke et al., 1997, Danen-van Oorschot, 1997a). Two, 3, 4 or 5 days after transfection the percentage of apoptotic apoptin-positive cells did not significantly exceed the percentage of apoptotic cells observed in the desmin-positive cultures, indicating that apoptin does not induce apoptosis in normal embryonic cell cultures. Transient transfection of transformed mouse/rat embryonic or baby rat kidney cells with the plasmid encoding apoptin proved that apoptin is able to induce apoptosis in these cells. The results of apoptin expression in "normal" embryonic rodent versus transformed rodent cells are shown in Figure 1.

These data show that apoptin fails to induce apoptosis in normal adult and embryonic rodent cells, but does induce apoptosis in the virally transformed derivatives, at least in cell culture conditions.

5

**Co-expression of SV40 large T antigen and apoptin results in apoptin-induced apoptosis in normal human diploid cells.**

We have examined the effect of expression of transforming genes on apoptin-induced apoptosis in normal human cells derived from healthy individuals. To that end, human VH10 diploid fibroblasts and FSK-1 diploid keratinocytes were transiently co-transfected with plasmid pCMV-VP3 encoding apoptin and either plasmid pSV40 encoding both SV40 large T and small T antigen, pR-s884 encoding large T antigen, pR-SVt encoding small T antigen, or the negative-control plasmid pCMV-neo. By indirect immunofluorescence, the cells were analysed for apoptin-induced apoptosis. Both normal VH10 and FSK-1 cells did not undergo apoptosis when apoptin was transfected with the control plasmid. The results showed, as expected, that expression of apoptin alone is not able to induce apoptosis in normal human diploid cells, confirming the data described by Danen-Van Oorschot (1997). However, normal diploid human fibroblasts and keratinocytes expressing both apoptin and SV40 large T antigen, alone or together with small T antigen, underwent apoptin-induced apoptosis (Figure 2). The rate of apoptosis induction was considerably increased in the presence of the viral transforming genes. Co-expression of the SV40 small T antigen with apoptin did not result in induction of apoptosis by apoptin. The transition of the normal cells, from apoptin-resistance to apoptin-susceptibility, can probably be explained by the fact that the apoptin protein translocates from a cytoplasmic localization to a nuclear localization. This transition becomes apparent already approximately 2 days after transfection of the SV40 plasmids (Fig. 3). One can conclude that an event takes place, in this example due to



expression of a transforming product from a DNA-tumor virus, which results in the translocation of apoptin from the cytoplasm to the nucleus, which is followed by induction of apoptosis.

5

**Co-expression of SV40 large T antigen and apoptin results in apoptin-induced apoptosis in normal rodent diploid cells.**

Next, we have examined the effect of co-expression of transforming genes and apoptin on apoptosis induction in  
10 normal mouse embryo fibroblasts (MEF) derived from p53 +/+ mice or from transgenic p53 -/- mice. Both types of transiently transfected MEFs co-expressing the transforming SV40 large T gene with or without small T antigen in combination with apoptin, underwent very fast apoptosis,  
15 whereas MEF expressing apoptin together with a control plasmid or with a plasmid encoding the non-transforming small T antigen, did not result in apoptin-induced apoptosis. The results are shown in Figure 4.

Immunofluorescence analysis also revealed that co-expression  
20 of apoptin and SV40 large T antigen resulted in the cellular translocation of apoptin. In the examined MEFs apoptin is situated in the cytoplasm. Upon SV40 large T antigen expression, apoptin enters the nucleus, followed by the induction of apoptosis. As comparison, the percentage of  
25 apoptin-positive transformed mouse cells is also given in Figure 5.

These results indicate that apoptin does not induce apoptosis in both p53 +/+ and p53 -/- mouse fibroblasts, but does so upon expression of a transforming protein.  
30 This information is important since it is known that "normal" p53 -/- cells are very susceptible to spontaneous transformation and easily progress to more highly transformed phenotypes. Loss of p53 alone, however, is not sufficient to create a transformed character. Furthermore, this finding  
35 shows that apoptin can induce apoptosis upon expression of a

transforming protein in other mammalian cells than human cells.

**Co-expression of SV40 large T antigen and apoptin results in  
5 apoptin-induced apoptosis in normal diploid cells derived  
from cancer-prone human individuals.**

By means of transient transfection and immunofluorescence we have also examined the effect of apoptin in the normal fibroblasts F9605 and GM1492 which are  
10 derived from individuals who show an increased cancer incidence due to a genetic defect. Apoptin is not able to induce apoptosis in normal diploid cells from these cancer-prone individuals. However, upon expression of SV40 large T antigen, apoptin induces apoptosis (Figure 6) after entering  
15 the nucleus (data not shown).

These data confirm that diploid cells, from hereditary cancer-prone syndromes are not susceptible to apoptin, whereas they become so when they express a transforming gene. Thus, diploid cells from such hereditary syndromes are  
20 identical to "normal" diploid human cells in this assay.

**The effect on induction of apoptosis of covalent linkage of a  
SV40 large T antigen nuclear localization signal to the  
apoptin protein.**

25 Next, we have examined whether expression of a chimeric protein consisting of apoptin and the nuclear localization signal of SV40 LT antigen (amino acids N-Proline-Proline-Lysine-Lysine-Lysine-Arginine-Lysine-Valine-C of SV40 large T antigen covalently linked to the N-terminus of apoptin)  
30 results in the induction of apoptosis in non- and transformed human cells. The chimeric protein is called NLS-apoptin. To that end, non-transformed VH10 human fibroblasts and transformed human osteosarcoma-derived Saos-2 cells (Danen-van Oorschot et al., 1997) were transfected with a plasmid  
35 encoding the chimeric protein NLS-apoptin. In both cell types, expression of NLS-apoptin resulted in the nuclear

localization of apoptin and induction of apoptosis.

Expression of the non-apoptotic protein Green Fluorescent Protein (GFP; (Pines, 1995) covalently linked to the NLS entered the nucleus, which did not result in the induction of apoptosis.

These data prove that a modified apoptin enabling its nuclear localization in a cell-transformed-independent manner, will be able to translocate into the nucleus, followed by induction of apoptosis.

A fusion product of the first N-terminal 69 amino acids of apoptin and the non-apoptotic GFP protein does not result in induction of apoptosis, and which coincides with the fact that this chimeric protein does not enter the nucleus (Noteborn and Pietersen, 1998). Now we have covalently linked the 8 amino acids of the NLS to the N-terminus of the apoptin fragment consisting of the amino acids 1-69 (NLS-apoptin/1-69). Transfection of both non-transformed VH10 cells and tumorigenic human cells (such as human osteosarcoma-derived Saos-2 cells) with a plasmid encoding the NLS-apoptin/1-69 resulted in the nuclear localization of the NLS-apoptin/1-69 followed by induction of apoptosis.

These data indicate that besides the C-terminal part of apoptin, also the N-terminal part (1-69 a.a.) does so, when it is translocated into the nucleus. In these experiments, as expected, the NLS-GFP was translocated into the nucleus but did not result in the induction of apoptosis.

**Normal diploid cells from cancer-prone individuals undergo apoptin-induced apoptosis after UV-radiation.**

We have also examined the effect of UV-irradiation on apoptosis induction by apoptin on diploid cells. Diploid fibroblasts derived from healthy persons (VH25) or from individuals with a cancer-prone syndrome (LF2675T cells from a Li Fraumeni Syndrome patient, and 401 cells from a Lynch Type 2 Syndrome patient) were transiently transfected with a plasmid encoding apoptin. Before transfection, part of the

cells was UV-radiated. As negative control, the cells were transfected with a plasmid encoding the protein desmin.

All 3 cell types, VH25, LF2675T and 401, did not reveal apoptin-induced apoptosis without UV irradiation. In combination with UV irradiation, however, the LF2675T and 401 cells, but not the VH25 cells, underwent apoptin-induced apoptosis very rapidly. Although we have no explanation for this phenomenon, it appeared that it correlates with another cellular property. Diploid cells from patients who are cancer-prone due to a germline mutation in a tumor suppressor gene, show an unexpected reaction to UV irradiation. When normal diploid fibroblasts are treated with UV or another DNA damaging agent, they react with a large variety of transient responses, including activation of signal transduction pathways, induction of expression of a variety of genes, inhibition of cellular DNA replication and activation of SOS-like phenomena such as Enhanced reactivation (ER) and Enhanced mutagenesis (EM). Abrahams et al. (1996) have found that normal diploid fibroblasts from patients with a hereditary cancer predisposition due to a germline mutation in a tumor suppressor gene, show the same responses to UV irradiation as cells from normal individuals, except for one response: Enhanced reactivation. The ER response in cells from these patients is much higher than in cells from normal individuals, hence these patient cells are called ERsuper (+). The molecular-biological basis of the ER phenomenon is still unclear. The detection of ER is a time-consuming approach, as it is based on the measurement of the (enhanced) survival of a UV-irradiated virus in UV-damaged (or X-ray damaged) cells, compared to the survival in non-damaged cells. An assay based on apoptin-induced apoptosis upon UV-radiation is considerably simpler and faster (see below). The fact that apoptin becomes active in cancer-prone cells upon UV-radiation makes it also possible to study the ER process. There is evidence to indicate that ER plays an important role in the process of cancer induction by DNA-damaging agents.

**Normal diploid cells from cancer-prone individuals undergo apoptin-induced apoptosis after X-ray treatment.**

Next, we have also examined the effect of X-ray treatment on the apoptosis induction by apoptin on human  
5 diploid cells. Diploid fibroblasts derived from healthy individuals (VH10) or from persons with a cancer-prone syndrome such as LF2675 and 401 cells were transfected with a plasmid encoding apoptin. Before, transfection, part of the cells was treated with X-rays (dose; 5 gray). As negative  
10 control, the cells were transfected with a plasmid encoding the protein desmin.

As expected, all analysed non-irradiated cells of the cell lines : VH10, LF2675 and 401, did not show apoptin-induced apoptosis. In combination with X-ray treatment,  
15 however, the cell lines derived from the cancer-prone individuals underwent apoptosis, whereas the ones derived from healthy persons did not. Five days after transfection, the majority of these X-ray-treated apoptin-positive cancer-prone cells had become apoptotic. The cells treated with X-  
20 rays and expressing the non-apoptotic agent desmin, did not undergo apoptin-induced apoptosis.

These results imply that treatment with X-rays, causing DNA-damage such as the above described UV-C treatment, results in the induction of apoptosis by apoptin in normal  
25 non-transformed human cells.

**Diagnostic assay for cancer-inducing genes based on apoptin-induced apoptosis.**

Danen-Van Oorschot et al. (1997a) have reported that the  
30 cellular localization of apoptin is different in tumorigenic/transformed human cells in comparison to the localization in normal non-transformed cells. Furthermore, accumulation of apoptin in the nucleus correlates with apoptosis induction, whereas cytoplasmic localization  
35 correlates with cell viability and normal proliferative capacity.

Based on the present report, we are able to develop a diagnostic assay for the identification of cancer-inducing and/or transforming agents or genes. A first type of assay consists of transfecting "normal" cells, for instance from  
5 human or rodent origin, with a plasmid encoding apoptin, or infecting the cells with viral vectors expressing apoptin, together with a plasmid encoding a putative transforming/cancer-inducing gene. Subsequently, the cells will be examined, (1) for the ability to undergo apoptosis by  
10 the apoptin gene and (2) for a shift in the localization of apoptin from the cytoplasm to the nucleus.

The intracellular localization of apoptin can be determined, using an immunofluorescence assay with monoclonal antibodies specific for apoptin, such as CVI-CAV-85.1. If the percentage  
15 of apoptosis in normal cells co-expressing apoptin and the putative transforming/cancer-inducing gene is significantly higher than in apoptin-positive control cells expressing a control plasmid, one can conclude that the analysed gene may indeed have transforming/cancer-inducing activity.

20 A second example of a diagnostic test is based on the treatment of cultured normal diploid cells with a putative carcinogenic agent. The agent can be added, for instance, to the culture medium for various lengths of time. Subsequently, the cells are transfected with a plasmid encoding apoptin or  
25 infected with a viral vector expressing apoptin. This approach can also be carried out by first transfecting /infecting the normal cells, and then treating the cells with the agent to be tested. The subsequent steps of the assay are the same as described for the first type of diagnostic assay.

30

#### **Diagnostic assay for cancer-proneness**

The data presented in this report allow us to develop an assay to determine whether an individual with an unknown cellular/genetic background, is cancer-prone compared to  
35 normal healthy persons. Normal diploid cells from a cancer-prone individual are insensitive to apoptin-induced

apoptosis, but become so after treatment with UV- or X-rays or another DNA damaging agent. Below, an example of such a diagnostic assay is described based on the effect of UV-irradiation. This assay can also be carried out with other  
5 mutagenic/carcinogenic agents.

Primary normal diploid cells are isolated from a skin biopsy of the individual to be tested and cultured in a suitable medium. Next, the cells are irradiated with UV and subsequently transfected with a plasmid encoding apoptin, or  
10 infected with a viral vector expressing apoptin, or the cells are first transfected/infected and then irradiated. In parallel, diploid cells from a normal healthy individual will be used as a control.

By using an indirect immunofluorescence assay based on  
15 apoptin-specific Mab's, the cells are analysed for the presence of apoptin in the nucleus and/or for undergoing apoptosis. If the percentage of cells undergoing apoptosis among the apoptin-positive UV-treated cells is significantly higher than the percentage of apoptosis in UV-treated cells  
20 of a normal individual, this will be strong evidence that the individual from whom the cells are isolated, will be cancer-prone.

#### **Use of apoptin proteins in pharmaceutical formulations for 25 anti-cancer therapy.**

On the basis of the above mentioned results one can also develop methods to apply apoptin in anti-cancer therapy, not as a gene (DNA) but as a protein. Apoptin is a comparatively small protein, which makes it feasible to introduce it into  
30 cells as a protein. (If fragments of the apoptin protein still have the desired apoptotic effect on cancer cells, we will use protein fragments instead of the intact protein). Our aim is to develop effective pharmaceutical formulations that ensure stability of the active component (= apoptin or a  
35 fragment thereof) and, if possible specificity for the tumor cell to be targeted.

The neoplasias that we hope to treat with suitable apoptin-containing formulations, both curative and preventive, include: hereditary forms of colorectal cancer (Familial adenomatous polyposis (APC) and hereditary non-polyposis colorectal cancer (HNPCC), cancer of the liver (or other  
5 organs that can be treated with perfusion techniques), leukemias and lymphomas (to be treated via the blood circulation), skin tumors and possibly lung tumors (via the respiratory tract).

10

**The construction and analyses of an expression-plasmid for the generation of transgenic-apoptin mice.**

The fact that we have now observed that apoptin fails to induce apoptosis in cultured murine embryonic fibroblasts,  
15 let us conclude that apoptin may also be expressed in the intact embryo and adult mice without causing toxicity, at least embryos of a not too early stage of embryonic development. We have chosen an expression system based on the murine H-2Kb transcription unit, which allows constitutive  
20 expression of foreign genes during embryogenesis and at adult stages in various organs (Drezen et al., 1992; Morello et al., 1986).

Therefore, we have constructed the expression plasmid p21EcoA-Vp3-Eco, which expresses apoptin under regulation of  
25 the murine H-2Kb promoter. Furthermore, the expression vector contains the other H-2Kb elements, which will allow expression of the apoptin gene. Figure 8 shows a schematic representation of the transgenic-apoptin expression vector p21EcoA-Vp3-Eco.

30 By means of transient transfections of transformed Saos-2 cells with the plasmid p21EcoA-Vp3-Eco we were able to prove that apoptin indeed could be expressed in the context of the H-2Kb sequences. Furthermore, the expressed apoptin resulted in the induction of apoptosis to a similar extent as apoptin  
35 expressed by means of the plasmid pCMV-VP3 (see figure 9).



These results imply that the used expression vector p21EcoA-Vp3-Eco expresses apoptin in such a way that transformed cells will undergo apoptosis.

## 5 The generation of transgenic-apoptin mice

In total 300 fertilized oocytes were micro-injected with a DNA fragment comprising of the H-2Kb transcription unit and the apoptin gene and transferred to 11 foster mice. In total we had gathered a progeny of 51 newborn mice.

10 By means of Southern-blot analysis (Southern, 1975) of BamHI- or XbaI-digested mouse-tail DNA using a 32P-labeled DNA fragment consisting of the complete VP3 gene, it was shown that the apoptin/H-2Kb unit was integrated within the genomic DNA of in total 7 founder (F0) mice. All transgenic-apoptin  
15 mice were looking healthy. For unknown reasons, however, 1 transgenic-apoptin mouse died at an age of 5-6 weeks.

The transgenic-apoptin mice were mated with FVB males or females. Tail DNA from the progeny was analysed for the presence of the apoptin gene using a polymerase chain  
20 reaction (PCR)-analysis using the primers P1 (5'-CTCTCCAACAACATACT-CCACCCGG-3') and P2 (CTTATACGCCTTTT-GCGGTTCGGG-3'). From all F0 mice, we have got 1 or more transgenic-apoptin F1-mice (Figure 10).

All mice of the F1-generation of the transgenic-apoptin  
25 animals were proven to be viable and thus far do not show any pathological defect, which might be correlated with the expression of apoptin.

By means of Northern-blot analysis (Noteborn et al., 1992)  
30 the expression of the apoptin gene could, as expected, be determined in various organs.

### Description of the figures

Figure 1 shows the apoptin-induced apoptosis activity in  
5 "normal" rodent embryo fibroblasts versus cells of  
transformed rodent cell lines. The cells were transiently  
transfected with pCMV-VP3 . Subsequently, the cells were  
fixed at several time intervals after transfection and  
10 analysed by indirect immunofluorescence using the apoptin-  
specific Mab 85.1. The percentage of apoptin-positive cells  
that stained abnormally with DAPI is given as relative  
measure for the induction of apoptosis.

Figure 2 shows the effect of SV40 large T antigen and/or  
15 small T antigen on apoptin-induced apoptosis in fibroblasts  
and keratinocytes from normal individuals. VH10 and FSK-1  
cells were transiently transfected with plasmid pCMV-VP3 and  
pCMV-neo or pSV40 expressing SV40 large T and small T  
antigen, pR-s884 expressing SV40 large T antigen, and pR-SVt  
20 expressing the SV40 small t antigen. Subsequently, the cells  
were fixed at several time intervals after transfection and  
analysed by indirect immunofluorescence using the apoptin-  
specific Mab 85.1. The percentage of apoptin-positive cells  
that stained abnormally with DAPI is given as relative  
25 measure for apoptosis.

Figure 3 shows the location of apoptin in normal human  
diploid cells expressing only apoptin or together with SV40  
large T antigen and/or small T antigen. The same cells  
30 analysed in Figure 2, regarding the induction of apoptosis  
were examined also for location of apoptin in the nucleus or  
cytoplasm. The percentage of apoptin-positive cells  
containing apoptin in the nucleus and still have not  
undergone apoptosis are given as relative measure of apoptin  
35 localization in the nucleus.

Figure 4 shows the effect of SV40 large T antigen and/or small T antigen on apoptin-induced apoptosis in mouse fibroblasts, which are derived from normal p53 +/+ mouse or from a transgenic p53 -/- knock-out mouse. The cells were transiently transfected with plasmid pCMV-VP3, expressing apoptin and the control plasmid pCMV-neo or with pSV40 expressing SV40 large T antigen, pR-s884 expressing large T antigen, and pR-SVt encoding small T antigen. Subsequently, the cells were fixed at several time intervals after transfection and analysed by indirect immunofluorescence using the apoptin-specific Mab 85.1. The percentage of apoptin-positive cells that stained abnormally with DAPI is given as relative measure for apoptosis

Figure 5 shows the location of apoptin in mouse embryonic p53 +/+ or p53 -/- fibroblasts expressing only apoptin or together with SV40 large T antigen and/or small T antigen. The same cells were analysed in Figure 4 regarding the induction of apoptosis were now also examined for location of apoptin in the nucleus or cytoplasm. The percentage of apoptin-positive cells containing apoptin in the nucleus and still being apoptotic are given as relative measure of apoptin localization in the nucleus.

Figure 6 shows the effect of SV40 large T antigen on apoptin-induced apoptosis activity in "normal" diploid human fibroblasts 9605 or G4905 derived from cancer-prone individuals. The cells were transiently transfected with pCMV-VP3 and pCMV-neo, or pSV40 expressing both large T and small T antigen of SV40, pR-s884 expressing large T antigen and pR-SVt expressing small T antigen. Subsequently, the cells were fixed at several time intervals after transfection and analysed by indirect immunofluorescence using the apoptin-specific Mab 85.1. The percentage of apoptin-positive cells that stained abnormally with DAPI is given as relative measure for apoptosis.

Figure 7 shows the effect of UV-irradiation on apoptin-induced apoptosis in "normal" diploid fibroblasts derived from normal healthy individuals versus cancer-prone patients. The cells were mock-treated or treated with UV-light and subsequently transiently transfected with pCMV-VP3 or with pCMV-des. Finally, the cells were fixed at several time intervals after transfection and analysed by indirect immunofluorescence using the apoptin-specific Mab 85.1. The percentage of apoptin-positive cells that stained abnormally with DAPI is given as relative measure for apoptosis.

Figure 8 shows a schematic representation of the transgenic-apoptin expression vector.

Figure 9 shows the apoptin-induced apoptosis activity in Saos-2 cells. The cells were transiently transfected with p21EcoA-Vp3-Eco, pCMV-VP3 (both expressing apoptin) or with pCMV-des, expressing the non-apoptotic protein desmin. Subsequently, the cells were fixed at several time intervals after transfection and analysed by indirect immunofluorescence using the apoptin-specific Mab 85.1. The percentage of apoptin-positive cells that stained abnormally with DAPI is given as a relative measure for the induction of apoptosis.

25

Figure 10  
Schematic representation of the pedigree of the VP3 (apoptin)-transgenic mice. The white-colored boxes are founder mice. The yellow- and green-colored boxes represent the progeny (F1) of the various apoptin-transgenic founders.

30

### References

1. Abrahams, P.J., Huitema, B.A., and Van der Eb, A.J. (1984). Enhanced reactivation and enhanced mutagenesis of herpes simplex virus in normal human and xeroderma pigmentosum cells. *Molecular Cellular Biology* 4, 2341-2346.
- 5 Abrahams, P.J., Houweling, A., and Van der Eb, A.J. (1992). High levels of enhanced reactivation of Herpes simplex virus in skin fibroblasts from various hereditary cancer-prone syndromes. *Cancer Research* 52, 53-57.
2. Abrahams, P.J., Houweling, A., Cornelissen-Steijger, P.D.M., Arwert, F., Menko, F.H., Pinedo, H.M., Terleth, C., and Van der Eb, A.J. (1996). Inheritance of Abnormal Expression of SOS-like Response in Xeroderma Pigmentosum and Hereditary Cancer-prone Syndromes. *Cancer Research*, 56, 2621-2625.
- 10 3. Bellamy, C.O.C., Malcomson, R.D.G., Harrison, D.J., and Wyllie, H. 1995. Cell death and disease: The biology and regulation of apoptosis. *Seminars on Cancer Biology* 6, 3-12.
4. Brinster, R.L., Chen, H.Y., and Trumbauer, M.E. (1981). Mouse oocytes transcribe injected *Xenopus* 5S RNA gene. *Science* 211, 396-398.
- 20 5. Danen-van Oorschot, A.A.A.M., Fischer, D., Grimbergen, J.M., Klein, B., Zhuang, S.-M., Falkenburg, J.H.F., Backendorf, C., Quax, P.H.A., Van der Eb, A.J., and Noteborn, M.H.M. (1997). Apoptin induces apoptosis in human transformed and malignant cells but not in normal cells. *Proceedings National Academy Sciences, USA*: 94, 5843-5847.
- 25 6. Danen-van Oorschot, A.A.A.M., Den Hollander, A., Takayama, S., Reed, J., Van der Eb, A.J., and Noteborn, M.H.M. (1997). BAG-1 inhibits p53-induced but not apoptin-induced apoptosis. *Apoptosis*, In Press.
- 30 7. Dinsart, C., Cornelis, J. J., Klein, B., van der Eb, A. J., and Rommelaere, J. (1984). Transfection with extracellularly UV-damaged DNA induces human and rat cells to

- express a mutator phenotype towards parvovirus H-1. *Molecular Cellular Biology* 4, 324-
8. Drezen, J.M., Nouvel, P., Babinet, C., and Morello, D. (1992). Different regulation of class I gene expression in the adult mouse and during development. *The Journal of Immunology* 149, 429-437.
9. De Ronde, A., et al. (1989). The SV40 small t antigen is essential for the morphological transformation of human fibroblasts. *Virology* 171, 260-263.
10. Duke, R.C., Ocjius, D.M., Young, J, D-E. (1996). Cell suicide in health and disease. *Scientific American* December 1996, 48-55.
11. Earnshaw, W.C., 1995. Nuclear changes in apoptosis. *Current Opinion in Cell Biology* 7, 337-343.
12. Fischer, D.F., Gibbs, S., Van de Putte, P., and Backendorf, C. (1996). Interdependent transcription control elements regulate the expression of the SPRR2A gene during keratinocyte terminal differentiation. *Molecular and Cellular Biology* 16, 5365-5374.
13. Graham, F.L. and Van der Eb, A.J. (1973). A new technique for the assay of infectivity of human adenovirus 5 DNA. *Virology* 52, 456-467.
14. Gruis, N.A. et al. (1995). Homozygotes for CDKN2(P16) germline mutation in Dutch familial melanoma kindreds. *Nature Genetics*, 10: 351-353, 1995.
15. Hockenberry, D.M. (1994). Bcl-2 in cancer, development and apoptosis. *Journal of Cell Science, Supplement* 18, 51-55.
16. Jacks, T. Lessons from the p53 mutant mouse (1996). *Journal of Cancer Research and Clinical Oncology* 122, 19-27.
17. Jacks, T. (1994). Tumor spectrum analysis in p53-mutant mice. *Current Biology* 4, 1-7.
18. Kerr, J.F.R., Winterford, C.M., and Harmon, B.V. (1994). Apoptosis: Its significance in cancer and cancer therapy. *Cancer* 73, 2013-2026.
19. Klein, B., Pastink, A., Odijk, H., Westerveld, A., and Van der Eb, A.J. (1990). Transformation and immortalization

- of diploid Xeroderma pigmentosum fibroblasts. *Experimental Cellular Research* 191, 256-262.
20. Levine, A.J. (1997). P53, the cellular gatekeeper for growth and division. *Cell* 88, 323-331.
- 5 21. Lowe, S.W., et al. (1994). Abrogation of oncogene-associated apoptosis allows transformation of p53-deficient cells. *Proceedings of the National Academy of Sciences, USA* 91, 2026-2030.
22. Maniatis, T., Fritsch, E.F., and Sambrook, J. (1982). *Molecular Cloning: A Laboratory Manual*. CSHL Press, New York, USA.
- 10 23. McBurney, M. W. et al. (1982). Isolation of male murine embryonal carcinoma cells and their chromosome replication patterns. *Developments in Biology* 89, 503- 508.
- 15 22. McDonnell T.J., Meyn, R.E., Robertson, L.E. (1995). Implications of apoptotic cell death regulation in cancer therapy. *Seminars in Cancer Biology* 6, 53-60.
23. Menke, A.L., Shvarts, A., Riteco, N., Van Ham, R.C.A., Van der Eb, A.J., and Jochemsen, A.G. (1997). Wilms' Tumor 1 (WT1) splice variants lacking the KTS induce apoptosis in p53-negative and p53-positive HepG2 cells. *Cancer Research* 57, 1353-1363.
- 20 24. Mellor, A.L., Golden, L., Weiss, E., Bullman, H., Hurst, Simpson, E., James, R.F., Townsend, A.R., Taylor, P.M., Schmidt, W., Ferluga, J., Leben, L., Santamaria, M., Atfield, G. Festenstein, H., Flavell, R.A. (1982). Expression of murine H-2Kb histocompatibility antigen in cells transformed with cloned H-2 genes. *Nature* 298, 529-534.
- 25 25. Morello, D., Moore, G., Salmon, A.M., Yaniv, M., and Babinet, C. (1986). Studies on the expression of an H-2K/human growth hormone fusion gene in giant transgenic mice. *The EMBO Journal* 5, 1877-1883.
- 30 26. Noteborn, M.H.M. and Pietersen, A.M. (1998). PCT application 98/00213 entitled Adenovirus vector.
- 35 27. Noteborn, M.H.M. (1996). PCT application WO 96/41191. Apoptin induces apoptosis in human transformed and malignant

cells but not in normal cells as essential characteristic for the development of an anti-tumor therapy.

28. Noteborn and De Boer, G.F. (1996). Patent USA/no. 030, 335.
- 5 29. Noteborn, M.H.M., De Boer, G.F., Van Roozelaar, D., Karreman, C., Kranenburg, O., Vos, J., Jeurissen, S., Zantema, A., Hoeben, R., Koch, G., Van Ormondt, H., and Van der Eb, A.J. (1991). Characterization of cloned chicken anemia virus DNA that contains all elements for the  
10 infectious replication cycle. Journal of Virology 65, 3131-3139.
30. Noteborn, M.H.M., Hoeben, R.C., and Pietersen, A. (1997). A gene delivery vehicle expressing the apoptosis-inducing proteins VP2 and/or apoptin. European Patent  
15 Application no. 97201121.7
31. Noteborn, M.H.M., Todd, D., Verschueren, C.A.J., De Gauw, H.W.F.M., Curran, W.L., Veldkamp, S., Douglas, A.J., McNulty, M.S., Van der Eb, A.J., and Koch, G. (1994). A single chicken anemia virus protein induces apoptosis.  
20 Journal of Virology 68, 346-351.
32. Noteborn, M.H.M., Kranenburg, O., Zantema, A., Koch, G., De Boer, G.F., and Van der Eb, A.J. (1992). Transcription of the chicken anemia virus (CAV) genome and synthesis of its 52-kDa protein. Gene 118, 267-271.
- 25 33. Paulovich, A.G., Toczyski, D., Hartwell, H. (1997). When checkpoints fail. Cell 88, 315-321.
34. Philips, B., and Rundell, K. (1988). Failure of semian virus 40 small t antigen to disorganize actin cables in nonpermissive cell lines. Journal of Virology 62, 768-775.
- 30 35. Pines, J. (1995). GFP in mammalian cells. Trends in Genetics 11, 326-327.
36. Ponc , M., Kempenaar, J.A., and De Kloet, E.R. (1981). Corticoids and cultured human epidermal keratinocytes: specific intra-cellular binding and clinical efficacy.  
35 Journal Investmental Dermatology 76, 211-214.
37. Reinwald, J.G. and Green, H. (1975). Cell 6, 331-343.



38. Sachs, L. and Lotem, J. (1993). Control of programmed cell death in normal and leukemia cells: New implications for therapy. *Blood* 82, 15-21.
39. Schrier, P. I., Bernards, R., Vaessen, R. T. M. J., Houweling, A. and Van der Eb, A. J. (1983). Expression of class I major histocompatibility antigens switched off by highly oncogenic adenovirus 12 in transformed rat cells. *Nature*, 305, 771-775.
40. Smits, P.H.M. et al. (1992). Modulation of the human papillomavirus type 16 induced transformation and transcription by deletion of loci on the short arm of human chromosome 11 can be mimicked by SV40 small t. *Virology* 190, 40-44.
41. Southern, E.M. (1975). Detection of specific sequences among DNA fragments separated by gel electrophoresis. *Journal of Molecular Biology* 98, 503-517.
42. Srivastava, S., Zou, Z., et al. (1990). Germline transmission of a mutated p53 gene in a cancer-prone family with Li-Fraumeni syndrome. *Nature* 348, 747-749.
43. Steller, H. (1995). Mechanisms and genes of cellular suicide. *Science* 267, 1445-1449.
44. Telford, W.G., King, L.E., Fraker, P.J. (1992). Comparative evaluation of several DNA binding dyes in the detection of apoptosis-associated chromatin degradation by flow cytometry. *Cytometry* 13, 137-143.
45. Teodoro, J.G. and Branton, P.E. (1997). Regulation of apoptosis by viral gene products. *Journal of Virology* 71, 1739-1746.
46. Thompson, C.B. (1995). Apoptosis in the pathogenesis and treatment of disease. *Science* 267, 1456-1462.
47. Van den Heuvel, S.J.L., Van Laar, T., Kast, W.M., Melief, C.J., Zantema, A., and Van der Eb, A.J. (1990). Association between the cellular p53 and the adenovirus 5 E1B-55kD proteins reduces the oncogenicity of Ad-transformed cells. *EMBO Journal* 9, 2621-2629.

48. Van Laar T., Steegenga wt et al. (1994). Bloom's Syndrome cells GM1492 lack detectable p53 protein but exhibit normal G1 cell-cycle arrest after UV irradiation. *Oncogene*. 9: 981-983
- 5 49. White, E. (1996). Life, death, and the pursuit of apoptosis. *Genes and development* 10, 1-15.
50. Wyllie, A.H. (1995). The genetic regulation of apoptosis. *Current Opinion in Genetics and Development* 5, 97-104.
- 10 51. Wyllie, A.H., Kerr, J.F.R., Currie, A.R. (1980). Cell death: The significance of apoptosis. *International Review of Cytology* 68, 251-306.
52. Zhuang, S.-M., Landegent, J.E., Verschueren, C.A.J., Falkenburg, J.H.F., Van Ormondt, H., Van der Eb, A.J.,
- 15 Noteborn, M.H.M. (1995). Apoptin, a protein encoded by chicken anemia virus, induces cell death in various human hematologic malignant cells in vitro. *Leukemia* 9 S1, 118-120.
53. Zhuang, S.-M., Shvarts, A., Van Ormondt, H., Jochemsen, A.-G., Van der Eb, A.J., Noteborn, M.H.M. (1995). Apoptin, a
- 20 protein derived from chicken anemia virus, induces a p53-independent apoptosis in human osteosarcoma cells. *Cancer Research* 55, 486-489.

CLAIMS

1. Method for determining the transforming capability of a possible transforming agent, comprising providing a non-transformed cell with inducible apoptin-like apoptotic activity, exposing said cell to said transforming agent and  
5 determining the localization of said apoptotic activity within said cell or determining the induction of apoptosis in said cell.
2. Method according to claim 1 whereby said apoptotic activity is provided by transducing said cell with a  
10 recombinant nucleic acid molecule encoding said activity.
3. Method according to claim 1 or 2, whereby said possible transforming agent is a proteinaceous substance.
4. Method according to claim 3 whereby said proteinaceous substance is co-expressed in said non-transformed cell with  
15 said apoptotic activity.
5. Method for determining the predisposition of a cell to become a tumor cell, by providing said cell with inducible apoptin-like apoptotic activity and subjecting said cell to relatively mild tumorigenic activity and determining  
20 apoptosis in said cell and/or determining the localization of said apoptotic activity in said cell.
6. Method according to claim 5 whereby said mild tumorigenic activity is UV-irradiation.
7. Method for determining the predisposition of a subject for  
25 hereditary types of cancer, comprising subjecting a sample of a relevant subset of cells of such a subject to a method according to claim 5 or 6.
8. Method for determining a gene mutation having oncogenic and/or transforming activity in a cell comprising subjecting  
30 said cell to a method according to claim 5 or 6.
9. Use of a nucleic acid encoding apoptin or a functional derivative or fragment thereof in the preparation of a

medicament for the prophylactic treatment of subsets of cells in a person, which subset of cells is cancer-prone.

10. Use according to claim 9 wherein said nucleic acid is provided in the form of a gene delivery vehicle.

5 11. Diagnostic test kit for carrying out a method according to anyone of claims 1-4, comprising a non-transformed cell transduced with a nucleic acid encoding apoptin or a functional derivative or fragment thereof.


10 12. Diagnostic test kit for carrying out a method according to anyone of claims 5-8, comprising a nucleic acid encoding apoptin or a functional derivative or fragment thereof capable of transducing a eukaryotic cell and capable of being expressed in such a cell.

15 13. Diagnostic test kit according to claim 12 comprising a means for subjecting a cell to a mild tumorigenic activity.

14. Method according to claim 4, whereby said proteinaceous substance is the large T-antigen of SV40 or a functional equivalent thereof.


## Figure 1

Apoptin-induced apoptosis in normal rodent embryo fibroblasts versus transformed rodent cell lines





## Figure 2

Effect of SV40 LT+st , LT or st antigen on  
apoptin-induced apoptosis in VH10 and FSK-1



## Figure 3



Location of apoptin in VH10 and FSK-1 expressing only apoptin or together with SV40 LT+st, LT or st antigen



4/10

## Figure 4


Effect of SV40 LT+st or LT antigen on apoptin-induced apoptosis in MEF(P53+/+) and MEF(p53-/-)







## Figure 5

Location of apoptin in MEF(p53+/+) or MEF(p53-/-) expressing only apoptin or together with SV40 LT+st or LT antigen



## Figure 6

Effect of SV40 LT+st or LT antigen on apoptin-induced apoptosis in normal diploid human fibroblasts derived from cancer-prone individuals



7/10

## Figure 7

Effect of UV-irradiation on apoptin-induced apoptosis in normal diploid fibroblasts derived from healthy versus cancer-prone individuals







Figure 8



9/10

Figure 9





Figure 10

## Overview VP3 transgenic mice



# INTERNATIONAL SEARCH REPORT

International Application No

PCT/NL 98/00457

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 G01N33/50 G01N33/574

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 G01N C07K A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 96 41191 A (AESCULAAP BV) 19 December 1996 cited in the application see page 6, line 22 - line 25	1-8, 14
X	see page 5, line 25 - line 30 see page 8, line 8 - line 20 ---	9, 10
X	DANEN-VAN OORSCHOT, A.A.A.M. ET AL.: "Apoptin induces apoptosis in human transformed and malignant cells but not in normal cells." PROC. NATL. ACAD. SCI. USA, vol. 94, May 1997, pages 5843-5847, XP002052232 cited in the application * first full paragraph of p. 5845, left column. * see the whole document ---	1-14
-/--		



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

\* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"Z" document member of the same patent family

Date of the actual completion of the international search

21 October 1998

Date of mailing of the international search report

02/11/1998

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax: (+31-70) 340-3016

Authorized officer

Hoekstra, S

# INTERNATIONAL SEARCH REPORT

International Application No

PCT/NL 98/00457

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	<p>NOTEBORN M ET AL: "CAV-derived apoptin induces apoptosis in human tumor cells in vitro"</p> <p>JOURNAL OF CONTROLLED RELEASE, vol. 41, no. 1, August 1996, page S10</p> <p>XP004037539</p> <p>see abstract</p>	11-13
X	<p>ZHUANG S -M ET AL: "APOPTIN, A PROTEIN ENCODED BY CHICKEN ANEMIA VIRUS, INDUCES CELL DEATH IN VARIOUS HUMAN HEMATOLOGIC MALIGNANT CELLS IN VITRO"</p> <p>LEUKEMIA, vol. 9, no. SUPPL. 01, October 1995, pages S118-S120, XP000602147</p> <p>cited in the application</p> <p>see the whole document</p>	11-13



# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/NL 98/00457

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9641191 A	19-12-1996	AU 5913696 A	30-12-1996
		CA 2221495 A	19-12-1996
		EP 0830604 A	25-03-1998
<hr/>			